

Subchondroplasty

<u>Postop</u>	<u>Goals</u>	<u>Precautions</u>	<u>Exercises</u>
Weeks 0-2 PT 1-2x/week HEP daily	Edema and pain control Maintain full extension Limit quad inhibition ROM As tolerated	PWB Avoid prolonged standing/walking No reciprocal stair ambulation	Quadriceps isometrics & re-education Patella mobilization Emphasize full passive extension P/A/AAROM flexion/extension Proprioception training Hip progressive resisted exercises
Weeks 2-6 PT 2-3x/week HEP daily	Full ROM Minimal swelling Normalize gait Ascend/Decent 8" stair Reciprocate stairs Return to normal ADLs	Progression to WBAT Avoid pain with therapeutic exercises and functional activities	Continue above Advance flexibility and proprioception Leg press Step up/step down program Elliptical
Weeks 6-12 PT 2-3x/week HEP daily	Full ROM Return to normal activity Run pain free Maximize strength/flexibility	WBAT Avoid pain with therapeutic exercises and functional activities Return to sport (MD directed)	Continue above Initiate running when able to descend 8" step without pain/deviation Plyometrics Agility and sport specific training Advance proprioception Advance LE strengthening

Defer to Limitations of Concomitant Procedure of Performed

Ryan R. Fader, M.D.

For more info: www.ryanfadermd.com

ryanfadermd@gmail.com

call 952-946-9777